

Determination of *When* an Outcome is Actualised in a Quantum Measurement using DNA-Photolyase System

Dipankar Home*, Rajagopal Chattopadhyaya**

*Department of Physics, Bose Institute, Calcutta 700009, India; dhom@boseinst.ernet.in

**Department of Biochemistry, Bose Institute, Calcutta 700054, raja@bic.boseinst.ernet.in

The biochemical attachment of photolyase to ultraviolet (uv) absorbed DNA molecules provides a method for registering whether a source has emitted photons. Here using laws of chemical kinetics and related experimental methods we argue that the *instant* after which this information becomes *discernible* can be *empirically determined* by retrodicting from relevant data *when* the photolyase binding to uv-absorbed DNA molecules has started occurring. Thus an empirically investigable twist is provided to the quantum measurement problem.

If a system is initially in a state $\psi(\psi = a\psi_1 + b\psi_2)$ which is a superposition of two states ψ_1 and ψ_2 that are eigenstates of a dynamical variable which is measured, a general characteristic of its interaction with a measuring device is that it results in a final state of the form

$$\Psi = a\psi_1\Phi_1 + b\psi_2\Phi_2 \quad (1)$$

where Φ_1 and Φ_2 are mutually orthogonal and *macroscopically distinguishable* states of the device. It is an ineluctable feature of linear unitary quantum mechanical treatment of *any* measurement process that the final state of system coupled to measuring apparatus has an entangled nonfactorisable form given by Eq.(1).

The much debated quantum measurement problem [1-8] stems from the *meaning* of a *pure state* wave function in quantum mechanics giving rise to an inherent *incompatibility* between a wave function of the form (1) and *actualisation* of the result of a measurement. A pure state in quantum me-

chanics means that each member (in this case, a system coupled to an apparatus) of an ensemble described by a pure state Ψ as given by Eq. (1) has the *same* wave function Ψ . Thus a pure state in quantum mechanics corresponds to a *homogeneous* ensemble whose members are *indistinguishable*. On the other hand, *definiteness* of individual outcomes requires that different outcomes are *distinguishable*. All measurements culminate in the final ensemble of systems coupled to apparatus which is essentially *heterogeneous*. A heterogeneous ensemble is, however, represented by a *mixed state* in quantum mechanics. Since within standard quantum mechanics under no unitary time evolution a pure state can evolve into a mixed state (see, for instance, [7] pp. 87-88), *how* to coherently accommodate within quantum mechanics the occurrence of distinguishable outcomes is thus an intriguing “paradox”. Not surprisingly, Weinberg [9] has called this “the most important puzzle in the interpretation of quantum mechanics.”

Resolution of the measurement problem requires either modifying or enlarging the standard framework of quantum mechanics in an appropriate way [10]. Schemes to this end are of two categories : (a) Approaches [11-15] which leave the mathematical formalism of quantum mechanics unmodified but introduce new elements into the conceptual framework. (b) Models [16-19] which modify the mathematical formalism (preserving the usual results of the standard formulation in their empirically verified domains) in order to provide a dynamical description of a measurement process in terms of an actual transition from a pure to a mixed state (the so-called “collapse of a wave function”).

A key ingredient in any of the above approaches is the notion that a measurement outcome has a definite *objective* (observer-independent) *reality* in the sense that once *actualised*, it remains “out there” so that the result can be inspected at *any* subsequent instant without perturbing it. This therefore requires a precise specification of *at what stage* a measurement result is recorded in a *stable* and *discernible* form. The schemes mentioned above use different *a priori* criteria for specifying this stage. An objective formulation of such a criterion amenable to empirical scrutiny is *necessary* for developing a satisfactory resolution of the measurement problem. On this question the present paper indicates a new direction of study by using biomolecular analogues of quantum measuring devices which promise to be of particular significance because biomolecules such as DNA “occupy a strategic position

between microscopic and macroscopic bodies” [20].

The specific example considered in this paper pertains to ultraviolet (uv) absorption by DNA molecules which develop new covalent bonds at certain sites (formation of uv-induced pyrimidine dimers) leading to a global readjustment of atomic positions. The displacement of an individual atom relative to its neighbouring atoms at such a damaged site is $\leq 2 \times 10^{-8} cm$ which cannot be directly observed by using an electron microscope. However, it becomes discernible by means of a curious biochemical property, viz. that if there are nearby *photolyase enzyme* molecules, some of them get preferentially attached to the damaged sites in the uv-absorbed DNA (uv-induced pyrimidine dimers in DNA have a chemical affinity towards photolyase which is a single polypeptide chain of 454 to 614 amino acids). That such an arrangement registering arrival of uv photons provides an instructive example of quantum measurement was earlier discussed by us [21]. We pointed out specific characteristics of this example in the context of the measurement problem. The central point is that since uv-damaged DNA molecule attached to photolyase records information about the emission of photons in a form that can be inspected at will any number of times at arbitrary instants, it is not essentially different from any readout device registering a measurement outcome.

The present paper lends a new twist to the above example by invoking suitably chosen features of *chemical kinetics* associated with the biochemical process of photolyase enzyme binding to uv-absorbed DNA molecules. We argue that it is possible to estimate retrodictively from suitable measurements *when* the enzyme attachment process began (say, at t_0). Note that t_0 signifies the instant from which information about the relevant measurement outcome (viz. that a source has emitted photons) is available in a *stable* and *discernible* form. That is, if one chooses to make a measurement at *any* $t \geq t_0$, some DNA molecules can always be found attached to photolyase thereby indicating that uv photons had been absorbed. However, as already mentioned, *no* physical significance can be associated with t_0 within standard quantum mechanics in the sense of signifying the onset of an actual transition from a pure to a mixed state. This example therefore helps to sharpen the quantum measurement problem by bringing it out in an empirically relevant form.

The measurement under consideration is as follows. Given a source which has a probability of emitting a pulse of, say, $10^9 \gamma$ photons (as explained later,

the reason for choosing this size of the pulse is for convenience in outlining a concrete feasible experimental arrangement), the measurement in question is designed to find out not only *whether* the source (left to itself for a certain time) has emitted such a pulse but also that *if* a pulse is emitted, one can find out from *which* instant onwards information about the pulse emission becomes available. In the first stage, the emitted pulse of $10^9\gamma$ photons interact with a pure CsI crystal proliferating into a pulse of 10^{15}uv photons . This output pulse is still *transient* which in itself does *not* constitute a stable record of measurement information. To convert this into a *stable* record, these uv photons need to be incident on a “detecting device” which in our case is an aqueous solution of DNA mixed with photolyase.

The combined state of source coupled with the “detecting device” mentioned above is given by (a particular form of Eq.(1))

$$\Psi = a\psi_e\Phi_A + b\psi_0\Phi_0 \quad (2)$$

where ψ_e, ψ_0 denote states of the source corresponding to emission and no emission and Φ_A, Φ_0 denote photolyase attached and unattached states of DNA respectively. Note that $|a|^2, |b|^2$ denote respectively the probabilities of the source emitting or not emitting photons within the specified time interval. While writing Eq.(2) we assume that the aqueous solution in this detecting arrangement comprises sufficient number of DNA and photolyase molecules so that in the event of the source emitting photons, an appreciable number of DNA molecules absorbing the photons get attached to photolyase. This binding occurs through a *time-evolving* chemical process. Here the role of photolyase is crucial for forming *macroscopically distinguishable* states Φ_A, Φ_0 . There are of course other methods as well for distinguishing between uv damaged and undamaged DNA (e.g., using nuclear magnetic resonance, x-ray crystallography or relevant chemical properties of other enzymes), but for our purpose the use of photolyase appears most convenient.

The crucial point is that (as we have argued earlier) Eq. (1) or Eq. (2) does *not* in itself account for the registration of a definite outcome. If one accepts the “completeness” of a wave function in specifying the state of an individual system, an *additional hypothesis* of the collapse of a wave function from a pure to a mixed state is required in order to explain the emergence of a measurement outcome. At which precise stage this putative transition occurs is an inherently contentious issue. In our specific example two different

viewpoints [21,22] are possible :

(a) The actual collapse of a pure state wave function (Eq.(2)) to a mixture of states $\psi_e\Phi_A, \psi_0\Phi_0$ occurs *only when* an observation is made by an external means to find out whether some DNA molecules have got attached to photolyase.

(b) Collapse of the wave function (Eq.(2)) into a mixture $\psi_e\Phi_A, \psi_0\Phi_0$ begins to take place at the instant t_0 when one or more DNA molecules in the solution *start* getting attached to photolyase, independent of whether being observed by external means. This point of view is motivated on the ground that at t_0 an initially homogeneous ensemble actually begins to split into two physically distinct subensembles comprising photolyase bound and unbound DNA respectively. Hence t_0 has an objective (observer-independent) significance that needs to be incorporated within the description of the collapse of a wave function.

We shall now indicate how it is possible to estimate from appropriately collected empirical data the instant t_0 photolyase binding to DNA begins. In other words, the *instant* from which information about the photon emission can be known (discernible by any external observer) is shown to be *empirically determinable*. Thus, in this specific context, the viewpoint (a) appears to be untenable. How to accommodate the viewpoint (b) within the various suggested forms of dynamical models of wave function collapse is a nontrivial issue [21,22] which needs to be studied in detail. Even for models addressing the measurement problem without using the idea of wave function collapse, this type of study involving biomolecular systems as measuring devices should provide useful constraints about specifying the stage at which a measurement outcome is recorded.

We now proceed to discuss specifics of the relevant experimental scheme. The rate of formation of uv-damaged *DNA-photolyase complex* (whose instantaneous concentration is denoted by $[PS]_t$) depends on the instantaneous concentrations of potential enzyme attachment sites in uv-damaged DNA denoted by $[S]_t$ as well as that of free or unbound photolyase denoted by $[P]_t$. This biochemical reaction follows the second order rate law

$$d[PS]_t/dt = k[P]_t[S]_t \quad (3)$$

where $[P]_t = P_0 - [PS]_t$ with P_0 being the initial concentration of photolyase,

$[S]_t = S_0 - [PS]_t$ where S_0 is the initial concentration of potential enzyme attachment sites produced in uv-damaged DNA and k is the second order rate constant. Phenomenological basis of Eq.(3) lies in random collision between uv-damaged DNA and photolyase arising from a diffusion process in the aqueous solution which ultimately result in specific chemical binding between the two. It is *experimentally verified* [23] that the uv-damaged DNA-photolyase complex formation obeys Eq. (3) and that for such a system the value of k ranges from $1.4 \times 10^6 M^{-1} s^{-1}$ to $4.2 \times 10^6 M^{-1} s^{-1}$.

An integrated form of Eq. (3) is given by

$$(P_0 - S_0)^{-1} \ln \{ [S_0(P_0 - [PS]_t)] / [P_0(S_0 - [PS]_t)] \} = k(t - t_0) \quad (4)$$

which expresses the time elapsed $(t - t_0)$ from the onset of the reaction at $t = t_0$ in terms of the initial concentrations S_0 , P_0 and the instantaneous concentration $[PS]_t$.

We shall now indicate the way Eq.(4) can be used for designing an *optimal* experimental arrangement that would permit us to estimate with controllable accuracy the time at which the attachment of photolyase to the uv-damaged DNA molecules begins (the instant $t = t_0$ of Eq.(4)). Typically, a certain fraction of the uv-exposed DNA molecules will be damaged by the actual absorption of uv photons. Among the photolyase molecules moving about randomly in the aqueous solution, the ones which are sufficiently close to the uv-damaged DNA molecules get attached to the specific sites. Then $[PS]_t$ increases gradually with time, reaching a plateau after a certain time following Eq.(4). For our purpose, the relevant parameters P_0 , S_0 need to be chosen such that *sufficiently long period* elapses before the plateau region sets in. This will enable withdrawing a number of aliquots from the sample and determining the fraction of photolyase attached to uv-damaged DNA sites at various time points.

The absorption of uv photons by DNA follows Beer's law given by

$$\ln \left(\frac{I}{I_0} \right) = A = \varepsilon c_m L \quad (5)$$

where I_0 is the incident intensity, I is the intensity emerging from the sample, A is known as the absorbance of the sample, ε is the extinction coefficient, c_m is the molar concentration and L is the pathlength traversed by the photons. For a given sample, both A and ε depend on the wavelength of incident radiation and nature of the solvent.

For the sake of concreteness, let us choose a $10^{-10}M$ concentration for the aqueous solution of synthetic DNA of 10 base pairs (1 base pair $\cong 600m_H$ where m_H is the mass of a hydrogen atom) containing a single potential pyrimidine dimer formation site (two adjacent thymines). For such a solution, $\varepsilon \cong 10^5 M^{-1}cm^{-1}$ corresponding to the uv wavelength $\sim 250-270$ nm; for a pathlength of $L = 10$ cm, A turns out to be 10^{-4} . Note that higher concentrations of DNA would make the attachment of photolyase too fast for our purpose.

It is operationally convenient to start with the DNA site concentration far in excess of the photolyase, i.e., $S_0 \gg P_0, [PS]_t$ whence the second order rate law effectively simplifies to a pseudo first order one. This implies that in Eq.(3) the product of k and $S_0 (\cong [S]_t)$ can be taken as the new first order rate constant. With a $10^{-10}M$ concentration for the 10 base pair DNA, one may choose a far less initial concentration of photolyase, say, $10^{-12}M$ whose changes due to binding with DNA can be determined with reasonable accuracy by using carbon radioisotope labelled photolyase [23].

We shall now estimate the number of uv photons required to completely convert all the adjacent thymines into pyrimidine dimers (potential photolyase attachment sites) in the 10 base pair DNA molecules. We consider that these molecules are in an aqueous solution contained within a size of $1mm \times 1mm \times 10$ cm (pathlength). Since $A = 10^{-4}$, this means that 0.023% of the incident uv photons are absorbed by such a solution. However, absorption of an uv photon does *not* necessarily lead to the formation of a pyrimidine dimer in a DNA molecule. Like any photoreaction, the formation of pyrimidine dimers depends on a quantum yield ϕ which is the ratio of the number of photons utilised in forming pyrimidine dimers to the number of photons actually absorbed. For a quantum yield $\phi = 0.015$ observed in the case of polynucleotides [24], it is calculated that 1.74×10^{15} incident uv photons are required to convert *all* the adjacent thymines into pyrimidine dimers in the solution whose concentration and size are indicated above. This means that if the source emits a pulse of $10^9 \gamma$ photons, the resulting 10^{15} uv photons are sufficient to convert a sizeable number of DNA molecules into dimers or potential photolyase attachment sites.

Now let us suppose that the above solution containing $10^{-10}M$ of 10 base pair DNA and $10^{-12}M$ of photolyase is left exposed (in conjunction with the CsI arrangement mentioned earlier) for a certain time to a source which has

a probability of emitting a pulse of $10^9\gamma$ photons within that time interval. Then to “read” the relevant outcome and to know “when” the outcome was registered, the above solution needs to be subjected to the following procedure. By withdrawing aliquots (random portions of the sample) from the solution at various times and running them through a polyacrylamide gel [25], two spatially separated radioactively labelled bands will result *if* the source has emitted photons : one band corresponding to the *unbound* photolyase ($P_0 - [PS]_t$) and the other to the photolyase *bound* to uv-damaged DNA ($[PS]_t$) which is *heavier*. The proportion of photolyase in these bands within polyacrylamide gel can be determined by measuring the radioactive counts from them. Then as indicated below, t_0 can be estimated by using Eq.(4) on the basis of such measurements for any pair of instants t_1, t_2 provided S_0 is much greater than P_0 (pseudo first order reaction). This condition is satisfied by the concentrations chosen so that Eq.(4) simplifies to

$$\ln[(P_0 - [PS]_t)/P_0] = -S_0k(t - t_0) \quad (6)$$

Using Eq.(6) for any pair of instants t_1, t_2 we get the following relation

$$\ln[(P_0 - [PS]_{t_1})/(P_0 - [PS]_{t_2})] = S_0k(t_2 - t_1) \quad (7)$$

Since left hand side of Eq.(7) is experimentally determined by the radioactive counts from photolyase bands within the polyacrylamide gel, (S_0k) can thus be calculated. This would enable one to estimate t_0 from Eq.(6) for either t_1 or t_2 . The half life associated with the pseudo first order process corresponding to Eq.(6) is given by $\ln 2/(S_0k)$. Substituting the values for S_0 and k mentioned earlier, this half life ranges from 57 to 83 minutes. Thus it is feasible to ensure a reasonably sufficient time before $[PS]_t$ reaches a plateau. One can therefore appropriately *increase* the accuracy in estimating t_0 by having more withdrawals and data at different times over the region where $[PS]_t$ varies with time significantly.

It should be noted that the operational significance of t, t_1, t_2 occurring in Eqs. (4), (6), (7) is that they refer to the instants of *applying* the withdrawn aliquots to the gel (*not* the instants of their withdrawal from the solution). This is because t, t_1, t_2 are the instants *till* which the photolyase attachment goes on before the separation between unbound and bound photolyase is made within the gel.

In the situation where the condition for the pseudo first order rate law is not

satisfied, one can still use Eq.(4) for computing t_0 but then the procedure is more complicated requiring data at three different instants. We also note that the time required for the formation of uv induced pyrimidine dimer in a DNA molecule is known to be exceedingly small, $\sim 10^{-14}s$ [26]; hence this period can be ignored in interpreting that t_0 signifies the instant from which the relevant outcome is registered.

To sum up, our preceding arguments show that the phenomenologically justified Rate Equation (whose various forms are Eqs. (3), (4), and (6)) entails *empirically determinable* t_0 . In the context of the quantum measurement problem, t_0 signifies the instant from which the outcome that a source has emitted photons *is knowable* in the sense that at *any* $t \geq t_0$, suitable observations can be made to “read” this outcome. In other words, t_0 signifies the *onset* of a *discernible heterogeneity* (mixed state) from an initial homogeneity (pure state). Thus t_0 has an objective factual significance which needs to be accommodated in a consistent way within the framework of quantum mechanics. To what extent the various models / interpretations of quantum mechanics can achieve this for such examples of quantum measurement calls for careful scrutiny. In particular, if a wave function is taken to provide a “complete” description of the state of a system, it becomes necessary to explain in detail the dynamical process by which the superposition of states given by Eq.(2) starts *reducing* to an actual mixed state of two terms ($\psi_e\phi_A$ and $\psi_0\phi_0$) in Eq. (2), *concomitant* with the onset of photolyase attachment to uv-absorbed DNA molecules.

Our scheme may be considered *analogous* to retrodictively inferring from suitable observations the *instant* at which a cat has expired upon the emission of photons in Schroedinger’s famous thought experiment [27], popularly known as the “cat paradox”. Our suggested new twist to the “cat paradox” makes it amenable to *controlled* experimental studies using biomolecular systems (in our example, photolyase attached DNA is analogous to the “dead” state of a cat in the context of “cat paradox”). Investigations along this direction have the potentiality to provide useful empirical clues and fresh insights into the quantum measurement problem, particularly because of *mesoscopic* sizes and masses of the biomolecules involved [28].

DH is grateful to Roger Penrose for raising insightful questions concerning our earlier work [Ref. 21] which helped in the formulation of the present work. DH also thanks John Corbett for stimulating discussions. RC acknowledges

research grant by BRNS, Govt. of India (4/5/95-R & D-II/703).

REFERENCES AND NOTES

1. A.J. Leggett, in *Quantum Implications*, B.J. Hiley, & D. Peat, Eds.(Routledge & Kegan Paul, London, 1987), pp. 85-104.
2. J.S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge University Press, Cambridge, 1987).
3. B. D'Espagnat, *Veiled Reality - An Analysis of Present -Day Quantum Mechanical Concepts* (Addison - Wesley, Reading, 1994).
4. J.T. Cushing, *Quantum Mechanics - Historical Contingency and the Copenhagen Hegemony* (University of Chicago Press, Chicago, 1994).
5. A. Whitaker, *Einstein, Bohr and the Quantum Dilemma* (Cambridge University Press, Cambridge, 1996).
6. M. Namiki, S. Pascazio, and H. Nakazato, *Decoherence and Quantum Measurement* (World Scientific, Singapore, 1997).
7. D. Home, *Conceptual Foundations of Quantum Physics - An Overview from Modern Perspectives* (Plenum Press, New York, 1997).
8. P. Mittelstaedt, *The Interpretation of Quantum Mechanics and the Measurement Process* (Cambridge University Press, Cambridge 1998).
9. S. Weinberg, *Dreams of a Final Theory*, (Vintage, London, 1993), p. 64.
10. Coupling of measuring apparatus with environmental degrees of freedom is often invoked to argue that coherence effects embodied in the pure state entangled form Eq.(1) are difficult to observe in practice (*effective decoherence*). The underlying contention is that the pure state (1) behaves *as if* it were a mixed state. Whether such “effective decoherence” between different outcomes is *sufficient* to ensure their *actual* distinguishability is highly debatable [1-4,7].
11. P. Holland, *The Quantum Theory of Motion* (Cambridge University

Press, Cambridge, 1993).

12. D. Bohm, and B.J. Hiley, *The Undivided Universe* (Routledge, London, 1993).

13. R. Griffiths, *J. Stat. Phys.* **36**, 219 (1984).

14. R. Omnes, *The Interpretation of Quantum Mechanics* (Princeton University Press, Princeton, 1994); *Understanding Quantum Mechanics* (Princeton University Press, Princeton, 1999).

15. N. Gisin, in *Fundamental Problems in Quantum Theory*, D.M. Greenberger, and A. Zeilinger, Eds. (Annals of New York Academy of Sciences, New York, 1995), pp. 524-533.

16. G.C. Ghirardi, A. Rimini, and T. Weber, *Phys. Rev. D* **34**, 470 (1986); P. Pearle, *Phys. Rev. A* **39**, 2277 (1989).

17. G.C. Ghirardi, R. Grassi and A. Rimini, *Phys. Rev. A* **42**, 1057 (1990); L. Diosi, *Phys. Rev. A* **40**, 1165 (1989).

18. G.C. Ghirardi, in *Fundamental Problems in Quantum Theory*, D.M. Greenberger, and A. Zeilinger. Eds. (Annals of New York Academy of Sciences, New York, 1995), pp. 506-523; G.C. Ghirardi, and R. Grassi, in *Bohmian Mechanics and Quantum Theory : An Appraisal*, J.T. Cushing, A. Fine, and S. Goldstein, Eds. (Kluwer, Dordrecht, 1996), pp. 353-377.

19. R. Penrose, *Gen. Rel. and Gravitation*, **28**, 581 (1996).

20. A. Shimony, in *Philosophical Consequences of Quantum Theory*, J.T. Cushing and E. McMullin, eds. (University of Notre Dame Press , Notre Dame , 1989), p. 61.

21. D. Home and R. Chattopadhyaya, *Phys. Rev. Lett.*, **76**, 2836 (1996).

22. P. Pearle and E. Squires, *Phys. Rev. Lett.*, **80**, 1348 (1998); see also Reply by D. Home and R. Chattopadhyaya, *Phys. Rev. Lett.*, **80**, 1348 (1998).

23. G.B. Sancar et al., *J. Biol. Chem.*, **262**, 478 (1987).

24. M.H. Patrick and R.D. Rahn, in *Photochemistry and Photobiology of Nucleic Acids*, S.Y. Wang, ed., (Academic Press, New York, 1976), Vol II,

pp. 35-145.

25. I. Husain and A. Sancar, *Nucl. Acids Res.*, **15**, 1109 (1987).

26. R.P. Wayne, in *Principles and Applications of Photochemistry*, (Oxford University Press, New York, 1998), pp. 30-32.

27. E. Schroedinger, *Naturwissenschaften*, **23**, 804 (1935).

28. For a quantitative idea about sizes and masses of the relevant biomolecules, see H.W. Park, S.T. Kim, A. Sancar and J. Deisenhofer, *Science*, **268**, 1866 (1995).